

Intracellular trehalose via transporter TRET1 as a method to cryoprotect CHO-K1 cells

メタデータ	言語: English 出版者: Elsevier 公開日: 2019-05-14 キーワード: trehalose, trehalose transporter 1, cryoprotectant, freezing temperature, trehalose concentration, CHO-K1 cell 作成者: 黄川田, 隆洋 メールアドレス: 所属:
URL	https://repository.naro.go.jp/records/2507

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



Intracellular trehalose via transporter TRET1 as a method to cryoprotect CHO-K1 cells

Tsutomu UCHIDA^{a*}, Maho FURUKAWA^a, Takahiro KIKAWADA^{b,c},
Kenji YAMAZAKI^a, and Kazutoshi GOHARA^a

^a Division of Applied Physics, Faculty of Engineering, Hokkaido University, N13 W8 Kita-ku, Sapporo,
Hokkaido 060-8628, Japan

^b Anhydrobiosis Research Group, Institute of Agrobiological Sciences, National Agriculture and Food Research
Organization, Ohwashi 1-2, Tsukuba, Ibaraki 305-8634, Japan

^c Graduate School of Frontier Sciences, The University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa, Chiba 277-8561,
Japan

* Corresponding author; e-mail: t-uchida@eng.hokudai.ac.jp; tel&fax: +81-11-706-6635

E-mail address: TU: t-uchida@eng.hokudai.ac.jp
MF: eje-w-_0w0_@eis.hokudai.ac.jp
TK: kikawada@affrc.go.jp
KY: k-yamazaki@eng.hokudai.ac.jp
KG: gohara@eng.hokudai.ac.jp

Abstract

Trehalose is a promising natural cryoprotectant, but its cryoprotective effect is limited due to difficulties in trans-membrane transport. Thus, expressing the trehalose transporter TRET1 on various mammalian cells may yield more trehalose applications. In this study, we ran comparative cryopreservation experiments between the TRET1-expressing cells and the wild-type cells under various trehalose concentrations in an extracellular medium and under various freezing temperatures. We show here that the intracellular trehalose significantly improves the viability of freeze-thawing cells. The optimum trehalose concentration is about 400 mM, and the optimum freezing temperature ranges between 140 and 240 K. We find a qualitative coincidence of the optimum freezing temperature with that from including 10 vol% dimethyl sulfoxide, a result that indicates that both intra- and extra-cellular trehalose principally work to inhibit ice crystal growth and thus protect the membrane and proteins from ice crystals during relatively slow freezing.

(144 words)

<Abbreviations>

TERT1, trehalose transporter 1; Me₂SO, dimethyl sulfoxide; SD, standard deviation

<Key words>

trehalose, trehalose transporter 1, cryoprotectant, freezing temperature, trehalose concentration, CHO-K1 cell

1 **Introduction**

2

3 Cultured cells from animals are widely used as alternative samples for pharmacological tests and as in-vitro
4 models in studies of principal biological mechanisms. The cells are usually incubated with a suitable culture
5 medium immediately after collection. However, due to an increased demand for such in-vitro models in tissue
6 engineering, cell transplantation, and genetic technologies, we need better long-term storage techniques for living
7 cells.

8

9 The traditional approach to such cell storage relies on cryopreservation methods that involve cryoprotectants such
10 as dimethyl sulfoxide (Me₂SO), glycerol, and ethylene glycol [13]. Recently, small carbohydrate sugars, such as
11 trehalose and sucrose, have been found to have an exceptional ability to stabilize and preserve cellular proteins
12 and membranes [3, 4, 12, 15, 22-24, 30]. In addition, trehalose inhibits ice-crystal growth [8, 25, 28]. However,
13 sugars such as glucose do not easily penetrate mammalian cells unless specific proteins are present in the cell
14 membrane to facilitate transport. Consequently, a number of methods have been explored to introduce non-native
15 sugars such as trehalose into mammalian cells. These include transfection [9], engineered pores [6, 7, 21],
16 activation of native channels [5], microinjection [6, 7], electropermeabilization [22], and endocytosis [10, 18].

17

18 Recent studies show that anhydrobiotic insect larvae use specific trehalose transporters during desiccation stress.
19 The trehalose transporter (TRET1) from the anhydrobiotic larvae of African chironomid, *Polypedilum*
20 *vanderplanki*, has been isolated and characterized [11, 14]. By stably expressing the TRET1 in mammalian cells,
21 extracellular trehalose was introduced into the cells with smaller stresses, and such cells significantly increased in
22 viability under partial desiccation [2].

23

24 Here, we investigate the cryopreserving effect of trehalose on mammalian cells expressing TRET1. Under various
25 trehalose concentrations in extracellular media and freeze temperature conditions, we measure the viabilities of
26 mammalian cells and obtain the optimum conditions. Comparing these results with those of Me₂SO-including
27 media, and with the physico-chemical properties of trehalose solution [28], we discuss the roles of trehalose as a
28 cellular cryoprotectant.

29

30 **Materials and methods**

31

32 ***Cell culture and transfection of CHO-K1 cells with trehalose transporters***

33

34 Gene-modified Chinese hamster ovary (Flp-In™-CHO) cells were purchased from Invitrogen (Thermo Fisher
35 Scientific, Waltham, MA, USA) and cultured in Nutrient Mixture F-12 Ham medium (Sigma-Aldrich, St. Louis,
36 MO, USA) supplemented with 10% fetal bovine serum (FBS, Thermo Fisher Scientific), 1% penicillin-
37 streptomycin (100 U/mL penicillin G and 100 µg/mL streptomycin, Thermo Fisher Scientific), 1% gluta max
38 (100x, Thermo Fisher Scientific), and 0.8% Hygromycin B solution (Nacalai Tesque, Kyoto, Japan). Cultures
39 were maintained at 310 K, and equilibrated with 5% CO₂-95% air in 75-cm² tissue culture flasks (AS ONE,
40 Osaka, Japan). Further details of the culture preparation are in a previous study [2].

41

42 The trehalose transporter (TRET1) expression vector (pcDNA5/FRT-PvTret1-AcGFP1; [supplementary Fig. S1](#))
43 for mammalian-cell expression was prepared and the TRET1 sequence was fused with a green fluorescent protein
44 (GFP) tag at the C-terminus by subcloning into the vector to generate a stable cell line. Stable cell lines were
45 generated using the Flip-in system (Thermo Fisher Scientific). Briefly, intact Flp-In™-CHO cells were transfected
46 with pcDNA5/FRT-PvTret1-AcGFP1 and pOG44 Flp-recombinase expression vector (Thermo Fisher Scientific)
47 using Fugene6 (Roche, Basel, Switzerland) and stably transfected cells were selected with hygromycin B, and
48 checked by the GFP expression on the cultured cell membrane with an epifluorescence microscope (IX71;
49 Olympus, Tokyo, Japan). We refer to this stable line as CHO-TRET1 cells. As a negative control, the
50 pcDNA5/FRT vector was transfected to Flp-In™-CHO cells, which is designated as CHO-vector cells.

51

52 ***Preparation of cell suspension for freezing***

53

54 When CHO-vector cell and CHO-TRET1 cells reached more than 80% confluence, the cells were dissociated
55 from the tissue culture flasks by trypsinization (0.05% trypsin-EDTA solution, Thermo Fisher Scientific) and
56 collected by centrifugation at 1000 rpm for 3 min. Then the cell-culture medium in the tissue culture flask was
57 removed completely using a Pasteur pipette with vacuum pump and immediately replaced with the culture media
58 that was supplemented with 0 ~ 1000 mM trehalose (purity 99.9%, endotoxin free, donated by Hayashibara). One
59 milliliter of the cell suspension (10⁵ cells/mL) was put into a 1.8-mL cryotube vial (Thermo Fischer Scientific)
60 and incubated for 6 h under the conditions described above for loading trehalose.

61

62 ***Cryopreservation protocol***

63

64 Following trehalose loading, both CHO-vector and CHO-TRET1 cells in solution were put into a freezer at
65 temperature settings of 263, 253, 243, 233, 213, and 198 K. Under these temperatures, the average rate of
66 temperature decrease for the samples (including 1-mL medium) V_f is about 5 K/min. Here we calculate V_f as the
67 temperature difference from room temperature to the freezing-state temperature divided by the time period prior
68 to the freezing start. To investigate temperatures down to below 198 K, we used the temperature gradient in a
69 partly filled liquid nitrogen dewar (Taylor-Wharton, Theodore, AL, USA). When about 770-mL liquid nitrogen
70 was put in the 5-L container, we inserted the vials. One vial was clipped at the lowest part of the vial holder (Shur-
71 Bend, St. Paul, MN, USA), which contacts the liquid nitrogen (77 K). The other four vials were clipped in upper
72 parts of the holder and thus held above the liquid nitrogen. Their temperatures were approximately 100, 140, 170,
73 and 200 K. Under these temperatures, V_f is estimated as ranging between 20 and 200 K/min. Each temperature
74 was measured by a thermocouple (T-type) on the side of each vial and recorded with a data logger (Graphtec,
75 Yokohama, Japan; type GL200A). After the temperature stabilizes to the environmental one, the sample is stored
76 at either the same temperature or in the deep freezer (198 K). As a preliminary survey showed no differences in
77 the viabilities whether stored at 198 K for one day or for two weeks, we fixed the storage period at one week (± 2
78 days).

79

80 To investigate the cryopreserving characteristics of trehalose, we also ran a similar test using 10 vol% Me₂SO
81 (Sigma), a popular trans-membrane type cryoprotectant. In this case, the sample was set in the dewar immediately
82 after replacing with the solution. We skipped the incubation procedure to inhibit the toxic effect of Me₂SO on the
83 cell activities.

84

85 ***Viability and growth assays***

86

87 The individual vials from each temperature conditions are then thawed in a 310-K water bath until all extracellular
88 ice is completely melted (about 5 min). During this process, the temperature increases by about 50 K/min. The
89 solution in the thawed cell suspension is then rinsed with culture media twice by centrifugation and the
90 supernatant solution removed. In this way, the cryoprotectants are largely removed from the culture media.

91

92 The survival rate of CHO-K1 cells was assessed using fluorescence dyes, calcein-AM and PI solutions (Cellstain;
93 Dojindo, Kumamoto, Japan), soon after the solution replacement (approximately 0.5-h after thawing). After
94 incubating for 20 min, we extracted 10 μ L from the cell suspension of live and dead cells and measured their
95 emissions at 515 and 620 nm on a hemocytometer (Waken counter, Waken B-tech, Kyoto, Japan). Fluorescence
96 images were captured using the epifluorescence microscope (IX71; Olympus) through an Olympus LCPLFL 20 \times
97 objective and collected with a charge-coupled device (DP70; Olympus). Cells of bright green fluorescence are
98 considered as alive, whereas bright red are scored as dead. Those displaying either both colors or no color were
99 used for estimating measurement uncertainties.

100

101 To determine the survival and growth efficiency, each CHO-cell suspension was replaced with the culture
102 medium and plated in a 24-well cell-culture plate (Nippon Genetics, Tokyo, Japan). The plated cells were then
103 incubated in a CO₂-incubator as described above for five days after thawing. The normality of the functions for
104 the living CHO-K1 cells was checked by observing the adhesion and growth of the cell, and qualitatively
105 comparing to the control CHO-K1 cells.

106

107 *Uncertainties of viability and statistical analysis.*

108

109 The viabilities are calculated as the number of living cells (stained by calcein-AM) normalized by total cell
110 populations (more than 500 cells). Considering the sampling variation, the staining error, and the human error of
111 counting the cells on a hemocytometer, the total uncertainty of any given data point is at most 8%. At least three
112 experiments were run per condition, so the resulting viability here is averaged at the indicated experimental
113 conditions \pm the standard deviation (SD) of all measured data. For the uncertainty of the data, we selected the
114 larger value between 8% of the mean value and the SD value.

115

116 For experiments on CHO-K1 cells, data were analyzed using two-way ANOVA with a Dunnett post-hoc test
117 (Prism, GraphPad Software) for at least 99% confidence ($p < 0.01$). All cryopreservation tests and 5-day growth
118 experiments were repeated at least three times for the independent cell lines.

119

120 **Results**

121

122 *Effect of trehalose concentration on cryopreservation of CHO-K1 cells*

123

124 We measured the effect of trehalose concentration in the medium on the TRET1-expressing CHO-K1 cells
125 viability and growth after freezing. **Figure 1** shows that after freezing and storing for 1 week at 193 K only a very
126 small fraction of both kinds of CHO-K1 cells survived in the absence of trehalose. The presence of extracellular
127 trehalose only slightly improves the viability of CHO-vector cells, but significantly improves the viability of
128 CHO-TRET1 cells. Moreover, the viability of the CHO-TRET1 increases with trehalose concentration.

129

130 To check the health of the cryopreserved cells, we cultured them at 310 K for one week and observed their
131 adhesion-proliferation processes. **Figure 2** shows micrographs of cryopreserved CHO-K1 cells under 500-mM
132 trehalose concentration for three-day culture. Adhesion of the surviving cells occurs after a one-day culture in
133 both types of cells (the left column of Fig. 2), although only few CHO-vector cells survive. After a two-day
134 culture, the CHO-TRET1 cells become almost confluent. Although the viability of CHO-vector cells is small,
135 those that survive can adhere and proliferate as normal cells. As these proliferation rates are almost the same as
136 the control, we consider that the surviving cells keep their normal functions after a freeze-thaw cycle.

137

138 As the cryopreserved cells were cultured in normal medium, the effect of intracellular trehalose on the health of
139 the cells is likely negligible. Our argument is based on a related experiment on oocyte: Kikawada et al. [11]
140 incubated oocyte in 105-mM trehalose buffer for three hours, then found that it took about two hours to eliminate
141 the intracellular trehalose from the oocyte. Given that CHO-K1 cells are smaller than oocytes, most of the
142 intracellular trehalose in our CHO-TRET1 cells should be removed within several hours. Hence, the cultivation
143 condition between CHO-vector and CHO-TRET1 cells are considered to be similar.

144

145 Now consider the effect of trehalose concentration on the viability of CHO-K1 cells. **Figure 3** shows the viability
146 of CHO-vector cells (gray diamonds) and CHO-TRET1 cells (black circles). The data roughly follow bell-shaped
147 curves, having relatively low viability at the lowest and highest trehalose concentrations. The maximum viabilities
148 are at 250 mM (0.099 ± 0.077) for CHO-vector and at 400 mM (0.806 ± 0.051) for CHO-TRET1 cells, which
149 indicates the viability is ten times larger in the latter case. Statistical analysis indicates a significant increase in
150 viability of CHO-TRET1 cells to that of CHO-vector cells ($p < 0.01$) at the trehalose concentration exceeding 200

151 mM. Consequently, we confirm that the trehalose loading into the cell with TRET1 significantly improves the
152 cryopreservation process.

153

154 *Effect of frozen temperature on cryopreservation of CHO-K1 cells*

155

156 Here we examine the temperature-dependence of the viability of the cryopreserved cells. Two concentrations are
157 considered: trehalose-free (0 mM) and the near-optimum concentration (500 mM). To better understand the
158 cryopreserving mechanism of the intracellular trehalose, we also used 10 vol% Me₂SO instead of trehalose.

159

160 For CHO-vector cells (Fig. 4a), the viabilities in both 0-mM and 500-mM trehalose concentrations are low (at
161 most 0.072 ± 0.060 for 0-mM trehalose at 145 K) at all temperatures. Conversely, the viabilities of CHO-TRET1
162 cells (Fig. 4b) are significantly higher for the 500-mM extracellular trehalose condition than those for the 0-mM
163 condition and those of CHO-vector cells at temperatures above 140 K ($p < 0.05$ at this temperature range). The
164 maximum viabilities occur at 213 K with a 500-mM concentration (0.740 ± 0.071). The results also show that the
165 intracellular trehalose improves the viability of the cryopreserved cells for temperatures at and above 140 K,
166 although the extracellular trehalose exhibits only a small cryopreservation effect.

167

168 Now compare the viabilities of CHO-K1 cells with trehalose to that with Me₂SO. Me₂SO is the most popular
169 trans-membrane type cryoprotectant, so we use it here as a positive control. For Me₂SO, more than 90% viability
170 occurs at temperatures between 130 and 230 K (marked as crosses in Figs. 4). This optimum temperature range
171 coincides well with that for CHO-TRET1 cells, which suggests that the intracellular trehalose cryoprotects via a
172 mechanism similar to that of Me₂SO. However, as the temperature exceeds 250 K, the Me₂SO loses its
173 cryoprotection ability for both CHO-vector and CHO-TRET1 cells, even though the intracellular trehalose retains
174 sufficient cryoprotecting effect. Thus, near the ice melting point, the intracellular trehalose appears to have
175 additional cryoprotecting roles than those of Me₂SO.

176

177 **Discussion**

178

179 In our freeze-thaw experiments, extracellular trehalose did not significantly affect cryopreservation of CHO-
180 vector cells. This result is consistent with that obtained for neurons with extracellular trehalose [16]. Conversely,

181 for the CHO-K1 cells loaded through TRET1, the intracellular trehalose did significantly increase the
182 cryopreservation. But what is the cryoprotecting mechanism of the intracellular trehalose? To address this
183 question, we examine how the viability was affected by the trehalose concentration and by the freezing
184 temperatures.

185

186 Consider first the cryopreservation of CHO-TRET1 cells at 193 K. The viability versus trehalose concentration in
187 the extracellular medium shows a bell-shaped curve in Fig. 3, peaking near 400 mM. This result is qualitatively
188 consistent with the viability of desiccation tolerance in CHO-K1 cells [2] and of cryopreservation of 3T3
189 fibroblasts and human keratinocytes [6], despite differences in experimental procedures and the findings of the
190 optimum condition. As the concentration of the intracellular trehalose is expected to be proportional to that of the
191 extracellular concentration [2, 11], this result suggests an optimum intracellular concentration for the
192 cryoprotecting effect of trehalose.

193

194 Concerning the minimum trehalose concentrations needed for sufficient cryoprotection, note that the viability of
195 CHO-TRET1 cells essentially vanishes for extracellular trehalose concentration below 200 mM. This behavior is
196 likely caused by having an insufficient amount of intracellular trehalose to prevent intracellular ice formation and
197 growth. Due to the relatively long loading time prior to freezing (6 hours), this intracellular-trehalose
198 concentration should be in equilibrium with the extracellular conditions. Hence, we argue that the minimum
199 extracellular trehalose concentration to show a sufficient cryoprotection effect is about 200 mM. By extrapolating
200 trehalose uptake data for CHO-TRET1 cells [2], this suggests a minimum intracellular trehalose concentration for
201 sufficient cryopreservation of about 27 nM/cell.

202

203 At the other extreme, the viability of the cryopreserved CHO-TRET1 cells decreases when the extracellular
204 trehalose concentration exceeds 500 mM. This result suggests that, even though trehalose is nontoxic for CHO-K1,
205 a higher intracellular trehalose concentration might weaken the cells. A possible reason for this result is a
206 weakening of CHO-TRET1 cells from hyperosmotic shock. As the Michaelis-Menten constant K_m for CHO-
207 TRET1 cells is 137 ± 87 mM [2], the maximum trehalose transport of TRET1 would be about 500 mM. Hence,
208 even when the extracellular trehalose concentration goes above 500 mM, the intracellular trehalose concentration
209 does not increase so much. If this effect occurs during incubation for the trehalose loading, the large amount of
210 extracellular trehalose would harm the cells via excessive dehydration. Another possible reason is that, as

211 trehalose has a high hydration effect, an oversupplied trehalose would reduce the intracellular water activities [28,
212 30], which may, for example, inhibit the intracellular material transport.

213

214 Second, consider the freezing-temperature dependence on the freeze-thawing viability of CHO-K1 cells. In Fig. 4,
215 we compare this viability for cells cryopreserved under two trehalose concentrations and one Me₂SO
216 concentration. High viabilities occur in both 500-mM-trehalose CHO-TRET1 cells and in the 10-vol%-Me₂SO
217 CHO-K1 (both vector and TRET1) cells at freezing temperatures between 140 and 240 K. This result suggests
218 that intracellular trehalose, in the optimum quantities, cryoprotects via a similar mechanism as Me₂SO.

219

220 About the mechanism for Me₂SO, we consider the solid-phase structures in Me₂SO solution. For such solution,
221 the glass transition temperature T_g' is 143 K, and the 3:1 compound, referred to here as the 'Me₂SO hydrate',
222 forms at temperatures between 143 and 210 K [17, 19]. In the 10 vol% (about 1.3 M) Me₂SO solution, the ice
223 melting point is about 260 K [17, 30]. This temperature is equivalent to the ice melting point in the culture
224 medium (without any additional cryoprotectant) [30]. Therefore, the Me₂SO hydrate, not the vitrification state, is
225 expected to form at the optimum cryoprotection temperature range (140 ~ 240 K). According to Rall et al. [19],
226 heterogeneous ice nucleation is inhibited under this optimum temperature condition and under a 10-vol% Me₂SO
227 concentration. Therefore, we argue that the optimum cryopreservation conditions in the present study are caused
228 by forming the Me₂SO hydrate, which then has a sufficient inhibition effect on ice nucleation and growth in the
229 cell. On the other hand, the glass-transition temperature T_g' of 500-mM trehalose solution should be a little lower
230 than 238 K, which is T_g' in 50 wt% (about 1.46 M) trehalose solution [1]. Thus, the high viabilities of 500-mM-
231 trehalose CHO-TRET1 cells at temperatures between 140 and 220 K may be caused by the vitrification of
232 intracellular solution, or by a sufficient inhibition effect of the intracellular trehalose on ice nucleation and growth.

233

234 Above 240 K, the viabilities of 10 vol%- Me₂SO CHO-K1 (both vector and TRET1) cells decrease drastically.
235 This temperature range is considered to be insufficient for the ice formation because of the reduction of the ice
236 freezing point down to about 260 K. For example, several samples with 10-vol% Me₂SO were not frozen during a
237 1-week-storage period at about 260 K (not shown in the plots). Therefore, the low viabilities of this temperature
238 range may occur either because the intracellular Me₂SO cannot inhibit the ice-crystal growth in the cell, or
239 because the cells are weakened by a toxic effect of Me₂SO [e.g., 29, 30] prior to the freezing.

240

241 The viability of 500 mM-trehalose CHO-TRET1 cells are as high as 0.5 at temperatures above 230 K. This high
242 viability in the trehalose-including conditions may be caused by a different mechanism than that for Me₂SO. As
243 the melting point of the medium including 500-mM trehalose is much higher than 230 K, the intracellular
244 trehalose would inhibit intracellular ice formation by reducing the activities of intracellular H₂O molecules [27,
245 28]. However, the cryoprotecting effect of trehalose occurs not only through inhibition of ice crystal growth [8, 25,
246 28] but also from a protection effect of cell membranes and the stabilization of proteins [3, 4, 12, 15, 22-24, 30]. If
247 the trehalose exists in both extra- and intra-cellular circumstances, the cell membrane may be sufficiently
248 protected both ways from the injury of ice crystal growth. The intracellular trehalose is expected not only to
249 protect the cell membrane and several organelles, but also to stabilize proteins in cytosol.

250

251 The slight decrease of viabilities near and above 230 K may be caused by the following mechanism. As the T_g' of
252 500-mM trehalose solution is just above 230 K, some of the intracellular water should be ice I_h [26] with further
253 growth inhibited by trehalose. The remaining intracellular water would include an amount of trehalose over the
254 optimum concentration, which reduces the activities of the intracellular H₂O molecules. This reduction would
255 reduce the viabilities of the cells rather than contribute to the cryoprotecting effect. The decrease in the viability at
256 high extracellular trehalose concentrations for CHO-TRET1 cells at 193 K (Fig. 3) supports this speculation.

257

258 However, below 140 K, the viabilities for both 500-mM trehalose CHO-TRET1 cells and 10 vol%- Me₂SO CHO-
259 K1 cells are low. This decrease is particularly sudden in the Me₂SO-administrated cells. As this critical
260 temperature coincides with the T_g' of 10 vol%- Me₂SO solution, the induction of a glassy state may be the cause.
261 This result differs from our expectation given that the cryoprotecting effect is caused by the vitrification of
262 intracellular condition. Rasmussen and MacKenzie [20] showed that vitrification required a 40 ~ 80 wt% Me₂SO
263 concentration for this particular cooling rate, which is much higher concentration than that in the present study.
264 Therefore, we suggest that the lower viabilities of 10 vol%- Me₂SO CHO-K1 cells at temperatures below 140 K
265 may be caused by intracellular freezing because the intracellular Me₂SO can neither induce vitrification nor form
266 the Me₂SO hydrate.

267

268 Consider the role of the freezing rate for the case of 500-mM trehalose CHO-TRET1 cells. The viabilities of 500-
269 mM trehalose CHO-TRET1 cells gradually decrease with temperature below 170 K. As the T_g' of 500-mM
270 trehalose solution is well above 140 K, no drastic changes, such as a phase change, could produce this decrease.

271 So, we consider instead the cooling rate V_f , which in these experiments varied with the freezing temperature. At
272 freezing temperatures above 180 K, V_f is below 10 K/min; At 140 K, it is about 20 K/min; and at 77 K, it is about
273 100 K/min. Based on these estimates, the viability of 500-mM trehalose CHO-TRET1 cells is high at V_f below 10
274 K/min, but decreases with increasing V_f above 10 K/min, and essentially vanishes when V_f exceeds 20 K/min.

275

276 The relation between V_f and the viability may be explained as follows: when V_f is large, the supercooling is large
277 (the ice nucleation occurs at lower temperatures). The larger the supercooling, the faster the ice-crystal growth,
278 and the greater difficulty for intracellular trehalose to inhibit the growth. According to Uchida et al. [25], ice
279 crystals formed in the 500-mM (~ 17 wt%) trehalose solution by quenching in liquid nitrogen were of order
280 several-microns across. This crystal size roughly equals that of CHO-K1 cells. When such an ice crystal forms in
281 the cell, the cell dies. To test this speculation, we plan to run additional experiments in which we control the
282 cooling rate.

283

284 We should also examine the melting rate. Uchida et al. [27, 28] showed that the existence of 10-vol% Me₂SO in
285 the culture medium reduced the melting point of ice below 265 K. As this temperature is below that of maximum
286 ice-grain growth, the frozen sample with Me₂SO should not have rapid ice recrystallization during melting.
287 Conversely, the ice melting point in the culture media with trehalose is about 270 K for a 500-mM trehalose
288 concentration [27, 28]. Thus, to understand the difference in viabilities of CHO-K1 cells with Me₂SO versus those
289 with trehalose, we should consider a possible effect of the rapid ice recrystallization during melting in the frozen
290 sample with trehalose. Future experimental studies including several additional parameterizations are needed.

291

292 **Conclusion**

293

294 We investigated the effect of intracellular trehalose transferred through TRET1 on the cryopreservation of CHO-
295 K1 cells. Our method involved changing either the trehalose concentration in the medium or the freeze-storage
296 temperature. We found that the intracellular trehalose improves the viability significantly. Cultivation of the
297 cryopreserved cells indicated that the surviving cells retain their functions, and therefore, that the trehalose
298 transferred via TRET1 minimizes the stresses on the cells. The optimum conditions for the cryopreservation of
299 CHO-TRET1 cells are (1) an extracellular trehalose concentration that exceeds 200 mM, with 250–500 mM being
300 optimal, and (2) a freezing temperature that exceeds 140 K, with 170–210 K being optimal.

301

302 To examine the cryoprotecting mechanism, we compared the temperature dependence of viabilities between the
303 optimum trehalose concentration and the most popular trans-membrane type cryoprotectant, Me₂SO. The results
304 indicated that the intracellular trehalose would not only inhibit the intracellular ice crystal growth, but also protect
305 the cell membrane and organelles as well as stabilize the proteins in cytosol. The latter effect becomes significant
306 at higher temperatures. An insufficient amount of intracellular trehalose, arising from a low extracellular trehalose
307 concentration, cannot provide cryoprotection. At the other extreme, too much intracellular trehalose may have
308 reduced the activities of intracellular water, thus lowering the measured viability. Even when the intracellular
309 trehalose concentration was near-optimal for cryopreservation, we found that the cooling rate should be below 10
310 K/min to prevent large supercooling conditions.

311

312 **Acknowledgments**

313 This work was partly supported financially by a Grant-in-Aid for Scientific Research from the Japan Society for
314 the Promotion of Science (Grant No. 26105009). Trehalose was donated by Hayashibara Co., Ltd. Cell cultures
315 and microscopic imaging are technically supported by Drs. D. Ito (Istitute Italiano di Tecnologia) and M.
316 Nagayama (Hokkaido University of Education). The cooling rate measurements were done by Mr. S. Hokonohara.
317 The authors acknowledge the discussion in the Joint Research Program of the Institute of Low Temperature
318 Science, Hokkaido University.

319

320 **References**

- 321 [1] C. Branca, S. Magazù, G. Maisano, P. Migliardo, F. Migliardo, G. Romeo, *α,α*-Trehalose/water solutions. 5.
322 Hydration and viscosity in dilute and semidilute disaccharide solutions, *J. Phys. Chem., B* 105 (2001) 10140-
323 10145.
- 324 [2] N. Chakraborty, M.A. Menze, H. Elmoazzen, H. Vu, M.L. Yarmush, S.C. Hand, M. Toner, Trehalose
325 transporter from African chironomid larvae improves desiccation tolerance of Chinese hamster ovary cells,
326 *Cryobiology* 64 (2012), 91-96.
- 327 [3] J.H. Crowe, L.M. Crowe, J.F. Carpenter, A.S. Rudolph, C.A. Wistrom, B.J. Spargo, T.J. Anchrodguy,
328 Interactions of sugars with membranes, *Biochim. Biophys. Acta* 947 (1988) 367-384.
- 329 [4] J.H. Crowe, L.M. Crowe, J.F., Preservation of mammalian cells – learning nature’s tricks, *Nat. Biotechnol.* 18
330 (2000) 145-146.

- 331 [5] G.D. Elliott, X.-H. Liu, J.L. Cusick, M. Menze, J. Vincent, T. Witt, S. Hand, M. Torner, Trehalose uptake
332 through P2X7 purinergic channels provides dehydration protection, *Cryobiology* 52 (2006) 114-127.
- 333 [6] A. Eroglu, M.J. Russo, R. Bieganski, A. Fowler, S. Cheley, H. Bayley, M. Toner, Intracellular trehalose
334 improves the survival of cryopreserved mammalian cells, *Nature Biotech.* 18 (2000) 163-167.
- 335 [7] A. Eroglu, J.A. Lawitts, M. Toner, T. Toth, Quantitative microinjection of trehalose into mouse oocytes and
336 zygotes, and its effect on development, *Cryobiology* 46 (2003) 121-134.
- 337 [8] T. Gonda, K. Sei, The inhibitory growth mechanism of saccharides on the growth of ice crystals from aqueous
338 solutions, *Progress in Crystal Growth and Characterization of Materials* 51 (2005) 70-80.
- 339 [9] N. Guo, I. Puhlev, D.R. Brown, J. Mansbridge, F. Levine, Trehalose expression confers desiccation tolerance
340 on human cells, *Nat. Biotechnol.* 18 (2000) 168-171.
- 341 [10] A. Hubel, T.B. Darr, Passive loading of trehalose into cells, *Cryobiology* 45 (3) (2002) 227.
- 342 [11] T. Kikawada, A. Saito, Y. Kanamori, Y. Nakahara, K. Iwata, D. Tanaka, M. Watanabe, T. Okuda, Trehalose
343 transporter 1, a facilitated and high-capacity trehalose transporter, allows exogeneous trehalose uptake into
344 cells, *PNAS* 104 (28) (2007) 11585-11590.
- 345 [12] C. Lambruschini, A. Relini, A. Ridi, L. Cordone, A. Gliozzi, Trehalose interacts with phospholipid polar
346 heads in Langmuir monolayers, *Langmuir* 16 (2000) 5467-5470.
- 347 [13] P. Mazur, Freezing of living cells: mechanisms and implications, *Am. J. Physiol.* 247 (1984) C125-C142.
- 348 [14] K. Mitsumasu, Y. Kanamori, M. Fujita, K. Iwata, D. Tanaka, S. Kikuta, M. Watanabe, R. Cornette, T. Okuda,
349 T. Kikawada, Enzymatic control of anhydrobiosis-related accumulation of trehalose in the sleeping
350 chironomid, *Polypedilum vanderplanki*, *FEBS J.* 277 (2010) 4215-4228.
- 351 [15] O. Miyawaki, M. Dozen, K. Hirota, Cooperative hydration effect causes thermal unfolding of proteins and
352 water activity plays a key role in protein stability in solutions, *J. Biosci. Bioeng.* 122 (2016) 203-207.
- 353 [16] J. Motomura, T. Uchida, M. Nagayama, K. Gohara, T. Taira, K. Shimizu, M. Sakai, Effects of additives and
354 cooling rates on cryopreservation process of rat cortical cells, in *Physics and Chemistry of Ice* (ed. W.F.
355 Kuhs), RSC Publishing, Cambridge, 2007, 409-416.
- 356 [17] S.S.N. Murthy, Some insight into the physical basis of the cryoprotective action of dimethyl sulfoxide and
357 ethylene glycol, *Cryobiology* 36 (1998) 84-96.
- 358 [18] A.E. Oliver, K. Jamil, J.H. Crowe, F. Tablin, Loading human mesenchymal stem cells with trehalose by fluid
359 phase endocytosis, *Cell Preserv. Technol.* 2 (1) (2004) 35-50.
- 360 [19] W.F. Rall, P. Mazur, J.J. McGrath, Depression of the ice-nucleation temperature of rapidly cooled mouse

- 361 embryos by glycerol and dimethyl sulphoxide, *Biophys. J.*, 41 (1983) 1-12.
- 362 [20] D.H. Rasmussen, A.P. MacKenzie, Phase diagram for the system water-dimethylsulphoxide, *Nature* 220
363 (1968) 1315-1317.
- 364 [21] M.J. Russo, H. Bayley, M. Toner, Reversible permeabilization of plasma membranes with an engineered
365 switchable pore, *Nature Biotechnol.* 15 (1997) 278-282.
- 366 [22] R. Shirakashi, C.M. Köstner, K.J. Müller, M. Kürschner, U. Zimmermann, V.L. Sukhrukov, Intracellular
367 delivery of trehalose into mammalian cells by electropermeabilization, *J. Membrane Biol.* 189 (2002) 45-54.
- 368 [23] A.K. Sum, R. Faller, J.J. de Pablo, Molecular simulation study of phospholipid bilayers and insights of the
369 interactions with disaccharides, *Biophys. J.* 85 (2003) 2830-2844.
- 370 [24] H. Takahashi, H. Ohmae, I. Hatta, Trehalose-induced destabilization of interdigitated gel phase in
371 dihexadecylphosphatidylcholine, *Biophys. J.* 73 (1997) 3030-3038.
- 372 [25] T. Uchida, M. Nagayama, T. Shibayama, K. Gohara, Morphological investigations of disaccharide molecules
373 for growth inhibition of ice crystals, *J. Crystal Growth* 299 (1) (2007) 125-135.
- 374 [26] T. Uchida, S. Takeya, Powder x-ray diffraction observations of ice crystals formed from disaccharide
375 solutions, *Phys. Chem. Chem. Phys.* 12 (45) (2010) 15034-15039.
- 376 [27] T. Uchida, M. Nagayama, K. Gohara, Trehalose solution viscosity at low temperatures measured by dynamic
377 light scattering method: Trehalose depresses molecular transportation for ice crystal growth, *J. Crystal
378 Growth* 311 (2009) 4747-4752.
- 379 [28] T. Uchida, S. Takeya, M. Nagayama, K. Gohara, Freezing properties of disaccharide solutions: inhibition of
380 hexagonal ice crystal growth and formation of cubic ice, in *Crystallization and materials science of modern
381 artificial and natural crystals* (eds. E. Borisenko, N. Kolesnikov), InTech, Rijeka, 2011, 203-224.
- 382 [29] T. Uchida, K. Miyamura, M. Nagayama, K. Gohara, Experimental investigations of cryopreservation
383 mechanisms on dispersed rat cardiac myocytes with dimethyl sulfoxide, *Cryobiology and Cryotechnology* 58
384 (1) (2012) 59-63 (in Japanese with English abstract).
- 385 [30] Z.-W. Yu, P.J. Quinn, Dimethyl sulphoxide: a review of its applications in cell biology, *Biosci. Rept.* 14
386 (1994) 259-281.
- 387

388 **Figure captions**

389

390 **Fig. 1.** Effect of trehalose concentration in the medium on the viability of cryopreserved cells. (a) CHO-vector
391 cells. (b) CHO-TRET1 cells. Samples stored at 193 K for 1 week. Red stain is PI, indicating dead cells, and green
392 stain is calcein-AM, indicating live cells. Horizontal distance across bottom is 2.2 mm.

393

394 **Fig. 2.** Micrographs of cells from the medium with 500-mM trehalose cryopreserved at 193 K for one week. (a)
395 CHO-TRET1 cells. (b) CHO-vector cells. After thawing in the bath at 310 K, the freezing medium was replaced
396 with culture media twice by centrifugation and removal of the supernatant solution. Each CHO-K1 cell suspension
397 was plated in a 24-well cell-culture plate. The plated cells were then incubated in a CO₂-incubator at 310 K for 5
398 days after thawing. The left column shows a 1-day culture after thawing, and the right column shows a 3-day
399 culture.

400

401 **Fig. 3.** Viabilities of cryopreserved CHO-vector cells (solid diamonds) and CHO-TRET1 cells (solid circles) at
402 various extracellular trehalose concentrations. The cells were incubated for six hours at 310 K prior to the sample
403 setting in the deep freezer (193 K). After 1-week storage, the frozen sample was thawed in the 310-K bath. Soon
404 after the thawing, the suspended cells were rinsed twice with PBS and incubated with fluorescence dyes. The
405 viability of each condition is the mean \pm SD (or measurement error) of epifluorescence microscopic measurements.
406 The asterisks mark cases with sufficient difference of viabilities between CHO-vector cells and CHO-TRET1 cells
407 at the same freeze-thaw conditions ($p < 0.01$).

408

409 **Fig. 4.** Effect of freezing temperature and trehalose transporter (TRET1) on the viability of cryopreserved cells.
410 (a) CHO-vector cells. (b) CHO-TRET1 cells. CHO-K1 cells suspended in the medium with trehalose at a
411 concentration (0 mM: squares, 500 mM: triangles) were set at the indicated temperature condition (77 – 263 K)
412 after a six-hour incubation at 310 K. The temperature dependence on viabilities with 10-vol% Me₂SO solution is
413 also plotted (marked as crosses). The viability of each condition is the mean \pm SD (or measurement error) of the
414 epifluorescence microscopic measurements.

1
2
3 Intracellular trehalose via transporter TRET1 as a method to cryoprotect CHO-K1 cells

4 (Figure files)

5
6 Tsutomu UCHIDA^{a*}, Maho FURUKAWA^a, Takahiro KIKAWADA^{b,c},
7 Kenji YAMAZAKI^a, and Kazutoshi GOHARA^a

8
9 ^a Division of Applied Physics, Faculty of Engineering, Hokkaido University, N13 W8 Kita-ku, Sapporo,
10 Hokkaido 060-8628, Japan

11 ^b Anhydrobiosis Research Group, Institute of Agrobiological Sciences, National Agriculture and Food Research
12 Organization, Ohwashi 1-2, Tsukuba, Ibaraki 305-8634, Japan

13 ^c Graduate School of Frontier Sciences, The University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa, Chiba 277-8561,
14 Japan

15 * Corresponding author; e-mail: t-uchida@eng.hokudai.ac.jp; tel&fax: +81-11-706-6635

16
17 E-mail address: TU: t-uchida@eng.hokudai.ac.jp

18 MF: eje-w-_0w0_@eis.hokudai.ac.jp

19 TK: kikawada@affrc.go.jp

20 KY: k-yamazaki@eng.hokudai.ac.jp

21 KG: gohara@eng.hokudai.ac.jp

22
23
24
25

Fig. 1




Fig. 1. Effect of trehalose concentration in the medium on the viability of cryopreserved cells. (a) CHO-vector cells. (b) CHO-TRET1 cells. Samples stored at 193 K for 1 week. Red stain is PI, indicating dead cells, and green stain is calcein-AM, indicating live cells. Horizontal distance across bottom is 2.2 mm.

Fig. 2




Fig. 2. Micrographs of cells from the medium with 500-mM trehalose cryopreserved at 193 K for one week. (a) CHO-TRET1 cells. (b) CHO-vector cells. After thawing in the bath at 310 K, the freezing medium was replaced with culture media twice by centrifugation and removal of the supernatant solution. Each CHO-K1 cell suspension was plated in a 24-well cell-culture plate. The plated cells were then incubated in a CO₂-incubator at 310 K for 5 days after thawing. The left column shows a 1-day culture after thawing, and the right column shows a 3-day culture.

Fig. 3




Fig. 3. Viabilities of cryopreserved CHO-vector cells (solid diamonds) and CHO-TRET1 cells (solid circles) at various extracellular trehalose concentrations. The cells were incubated for six hours at 310 K prior to the sample setting in the deep freezer (193 K). After 1-week storage, the frozen sample was thawed in the 310-K bath. Soon after the thawing, the suspended cells were rinsed twice with PBS and incubated with fluorescence dyes. The viability of each condition is the mean \pm SD (or measurement error) of epifluorescence microscopic measurements. The asterisks mark cases with sufficient difference of viabilities between CHO-vector cells and CHO-TRET1 cells at the same freeze-thaw conditions ($p < 0.01$).

Fig. 4




Fig. 4. Effect of freezing temperature and trehalose transporter (TRET1) on the viability of cryopreserved cells. (a) CHO-vector cells. (b) CHO-TRET1 cells. CHO-K1 cells suspended in the medium with trehalose at a concentration (0 mM: squares, 500 mM: triangles) were set at the indicated temperature condition (77 – 263 K) after a six-hour incubation at 310 K. The temperature dependence on viabilities with 10-vol% Me₂SO solution is also plotted (marked as crosses). The viability of each condition is the mean \pm SD (or measurement error) of the epifluorescence microscopic measurements.