

山形, 鳥取, 兵庫, 京都, 佐賀, 福岡, 大分, 長野県における マメ科植物遺伝資源の多様性保全 2007 年

友岡 憲彦¹⁾・加賀 秋人¹⁾・伊勢村 武久¹⁾・黒田 洋輔¹⁾・Asta Taman²⁾・
松島 憲一³⁾・根本 和洋³⁾・Duncan A. Vaughan¹⁾

1) 農業生物資源研究所・ジーンバンク

2) ブータン王国・王立生物多様性研究センター

3) 信州大学

Collection and Conservation of Leguminous Crops and Their Wild Relatives in Japan, 2007

Norihiko TOMOOKA¹⁾, Akito KAGA¹⁾, Takehisa ISEMURA¹⁾,
Yosuke KURODA¹⁾, Asta TAMAN²⁾, Ken-ichi MATSUSHIMA³⁾,
Kazuhiro NEMOTO³⁾ and Duncan A. VAUGHAN¹⁾

1) National Institute of Agrobiological Sciences, Kannondai 2-1-2, Tsukuba, Ibaraki, 305-8602, Japan

2) National Biodiversity Center, Sarbitan, Thimphu, Bhutan

3) Shinshu University, Minami-Minowa 8304, Kami-Ina, Nagano, 399-4511, Japan

Summary

Legume plant genetic resources were the target for exploration in Yamagata prefecture from September 24 to 25, Tottori, Hyogo, Kyoto, Saga, Fukuoka, Oita prefectures from October 22 to 25, and Nagano prefecture from November 5 to 6. As a result, 60 accessions of leguminous plants consist of 2 of *Glycine max*, 28 of *Glycine soja*, 7 of *Vigna angularis* var. *angularis*, 22 of *Vigna angularis* var. *nippensis*, 1 of *Vigna unguiculata* were collected. An invited Bhutanese scientist, Ms. T. Asta joined the survey based on the MOU between NIAS and National Biodiversity Center, Bhutan.

Introduction

In order to conserve genetic diversity of Japanese leguminous crops and their wild relatives, 3 exploration trips were made in 2007. We focused on wild *Glycine* (soybean) and wild *Vigna* (azuki bean) genetic resources.

Methods

We surveyed Yamagata prefecture from September 24 to 25, Tottori, Hyogo, Kyoto prefectures from October 22 to 23, Saga, Fukuoka, Oita prefectures from October 24 to 25, and Nagano prefecture from November 5 to 6 by car as shown in Fig. 1. Seeds, herbarium specimens and root nodules (if available) were collected. Information on collection sites including village name, altitude, latitude, longitude, habitat, cultural practices and other ecological data of the collection sites were recorded as passport data (Table 2).

Results and Discussion

A total of 60 legume accessions including the genus *Vigna* and *Glycine*, consist of 2 of *Glycine max*, 28 of *Glycine soja*, 7 of *Vigna angularis* var. *angularis*, 22 of *Vigna angularis* var. *nippensis*, 1 of *Vigna unguiculata* were collected. (Table 1 & 2).




Fig. 1. A map of wild legume genetic resources collection sites in 2007.

(September 24 to 25, Yamagata)

In Yamagata prefecture, 15 accessions of wild soybean (*G. soja*), 4 of wild azuki bean (*V. angularis* var. *nipponensis*, including 2 intermediate populations), and 1 of cultivated azuki bean (*V. angularis* var. *angularis*) were collected. Most of wild soybean populations are still in the early maturing stage and we could not collect mature seeds from some of the populations. In contrast, it was a good season to collect mature seeds from wild azuki bean populations. A wild soybean individual with umbrella like arrangement of pods, which has an erect main stem, was found in 2007-03 population (Photos 1 & 2).

An intermediate morphological wild soybean individual which is thought to be a hybrid between cultivated and wild plants was found at 2007-08 site (Urushizone, Sakata, Fig. 2, Photos 3 & 4). The intermediate individual was growing between the river levee and soybean field and had 6 lateral branches and 58 pods. There were soybean fields in the vicinity. The distance between the intermediate plant and the nearest soybean field was about 10 m. We collected 39 individual leaf samples from this wild soybean population at 1 m intervals from both sides of the intermediate plant. We also collected leaf samples from soybean fields. We plan to analyze the genetic structure of this population in detail.

At 2007-09 (Yamadate) and 2007-10 (Narahashi) sites, intermediate (weedy) azuki bean populations were found. At Yamadate site, several weedy azuki bean plants were growing between the river and a soybean field. They showed indeterminate growth habit. One of these plants had small red seeds. Despite the small seed size this plant may have been an escaped plant from cultivation and was collected separately as 2007-09C. At Narahashi site, weedy azuki bean plants were growing in and around a fallow paddy field. Seed and pod sizes were significantly larger than typical wild azuki bean. Plant type variation (determinate and semi-indeterminate) was observed within the population.




Fig. 2. A site where an intermediate morphological wild soybean individual was found (Urushizone, Sakata city, Yamagata prefecture). ○ indicates an intermediate soybean plant. ■ indicates wild soybean individuals from which leaf samples for DNA analysis were collected.

(October 22 to 23, Tottori, Hyogo, Kyoto)

In Tottori, Hyogo and Kyoto prefectures, 2 accessions of soybean, 6 of wild soybean, 6 of azuki bean, and 10 of wild azuki bean (including 2 intermediate accessions) were collected. Most of wild soybean populations are in the maturity stage, while most of wild azuki bean populations are in the past maturity stage. In the cultivated azuki bean fields in Kyotango city, Kyoto prefecture, one individual with black pods (2007Kyoto22-C) was found among white pod cultivars. This individual is thought to be a hybrid derivative with wild azuki bean. At this site, wild azuki bean and wild soybean also grow sympatrically (Photo 5). Two accessions of intermediate weedy azuki bean accessions were identified in a population (2007Hyogo24-A, Nakataeda, Tanba-Ichijima) based on the seed color difference (black mottled and pale brown, Photo 6).

(October 24 to 25, Saga, Fukuoka, Ooita)

In Saga, Fukuoka and Ooita prefectures, 3 accessions of wild soybean, 5 of wild azuki bean (including 1 intermediate weedy accession) and 1 of cowpea (*Vigna unguiculata*) were collected. In a population of intermediate weedy azuki bean (2007Ooita25, Uchikawano, Hita city), plants with green stem were growing on the road side slope and had experienced cutting (clearing by human) at least once. Wild soybean individuals were growing abundantly on the ground of sand digging place of mountain (2007Fukuoka27, Yamae, Chikushino city). Wild soybean populations are frequently found in this type of newly opened ground of sand digging place on mountain. In a similar habitat at Saga prefecture (2007Saga28, Sonobe, Kiyama), wild soybean and wild azuki bean were found growing sympatrically. At site 2007Saga29, Kase town, both wild soybean and wild azuki bean population were found growing. This is a paddy field area on the reclaimed land from the sea of Ariake.

(November 5 to 6, Nagano)

In Nagano prefecture, Ina and Shimo-Ina areas were explored. Four accessions of wild soybean and 3 accessions of wild azuki bean (including 2 intermediate weedy populations) were collected. Most of wild populations (soybean and azuki bean) were at the past maturity stage. A wild soybean population was found growing beside experimental field of the Shinshu University (2007N2). Two intermediate weedy azuki populations were found at Sankouchi, Yasuoka village (2007N3) and at Nakahira, Iida city (2007N7, Photo 8). The weedy azuki bean plants had green stems at both sites. However, weedy plants at Sankouchi showed twining growth, while weedy plants at Nakahira showed erect growth habit (Photo 8).

和文摘要

本報告は、2007年9月24～25日に山形で、10月22～23日に鳥取、兵庫、京都で、10月24～25日に佐賀、福岡、大分で、11月5～6日に長野で行ったマメ科植物遺伝資源探索の報告である。結果として60点のマメ科植物遺伝資源を保存した。その内訳は、ダイズ2点、ツルマメ28点、アズキ7点、ヤブツルアズキ22点、ササゲ1点である。山形県酒田市漆曾根（2007-08）のツルマメ集団の中に、栽培ダイズとの交雑後代と考えられる中間体を1個体発見した。ダイズの中間体は、この1ヶ所で確認できただけであるが、アズキの中間体は、調査したすべての地域で高頻度に確認できた。山形以外の探索には、ブータン王国・王立生物多様性研究センターとの研究協定に基づいて招聘したT. Astaジーンバンクマネージャーも参加した。

Table 1. Species and number of accessions collected in each region

日本各地の探索で収集した植物種と収集点数

| Species 種名 | No. of accessions collected in each region | | | | Total 合計 |
|---|--|-----------------------------------|--------------|----------------|-------------|
| | Saga, Fukuoka, Ooita 佐賀・福岡・大分 | Tottori, Hyogo, Kyoto 鳥取・兵庫・京都 | Nagano 長野 | Yamagata 山形 | |
| <i>Glycine max</i> ダイズ | - | 2 | - | - | 2 |
| <i>Glycine soja</i> ツルマメ | 3 | 6 | 4 | 15 | 28 |
| <i>Vigna angularis</i> var. <i>angularis</i> アズキ | - | 6 | - | 1 | 7 |
| <i>Vigna angularis</i> var. <i>nipponensis</i> ヤブツルアズキ | 5 (1)* | 10 (2)* | 3 (2)* | 4 (2)* | 22 |
| <i>Vigna unguiculata</i> ササゲ | 1 | - | - | - | 1 |
| Total 合計 | 9 | 24 | 7 | 20 | 60 |

* number in the parenthesis indicates number of intermediate weedy accessions

Table 2. A passport data of collected materials in Japan

収集品のパスポートデータ

| Coll. Date | Coll. No. | JP No. | Species name | Status | Collection Site (in Japanese) | Vill., City, Pref. | Latitude | Longitude |
|------------|--------------------------------------|--------|--|------------|----------------------------------|---|-------------|---------------|
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-01 | 233166 | <i>Glycine soja</i> | wild | 山形県鶴岡市播磨 | Harima, Tsuruoka-shi, Yamagata | N38-46-10.9 | E139-49-54.9 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-02 | 233167 | <i>Glycine soja</i> | wild | 山形県東田川郡三川町猪子 | Inoko, Mikawa-machi, Higashi Tagawa-gun, Yamagata | N38-49-23.6 | E139-50-24.2 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-03 | 233168 | <i>Glycine soja</i> | wild | 山形県酒田市坂野辺新田 | Sakanobeshinden, Sakata-shi, Yamagata | N38-53-29.2 | E139-49-56.8 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-04 | 233169 | <i>Glycine soja</i> | wild | 山形県酒田市藤塚 | Fujitsuka, Sakata-shi, Yamagata | N38-57-37.1 | E139-51-26.0 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-05 | 233170 | <i>Glycine soja</i> | wild | 山形県館海郡遊佐町吉出 | Yoshide, Yuza-machi, Akumi-gun, Yamagata | N39-01-44.1 | E139-54-28.4 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-06 | 233171 | <i>Glycine soja</i> | wild | 山形県飽海郡遊佐町吉出 | Yoshida, Yuza-machi, Akumi-gun, Yamagata | N39-01-05.3 | E139-56-18.4 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-07 | 233172 | <i>Glycine soja</i> | wild | 山形県酒田市下黒川 | Shimokurokawa, Sakata-shi, Yamagata | N38-59-42.1 | E139-58-23.3 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-08A | 233173 | <i>Glycine soja</i> | wild | 山形県酒田市漆曾根 | Urushizone, Sakata-shi, Yamagata | N38-55-30.4 | E139-53-08.4 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-08B | 233622 | <i>Glycine soja</i> | wild | 山形県酒田市漆曾根 | Urushizone, Sakata-shi, Yamagata | N38-55-30.4 | E139-53-08.4 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-09A | 233174 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 山形県酒田市山樋 | Yamadate, Sakata-shi, Yamagata | N38-54-09.7 | E139-56-09.7 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-09B | 233175 | <i>Glycine soja</i> | wild | 山形県酒田市山樋 | Yamadate, Sakata-shi, Yamagata | N38-54-09.7 | E139-56-09.7 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-09C | 233623 | <i>Vigna angularis</i> var. <i>angularis</i> | escape | 山形県酒田市山樋 | Yamadate, Sakata-shi, Yamagata | N38-54-09.7 | E139-56-09.7 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-10 ① | 233176 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 山形県酒田市橋橋 | Narahashi, Sakata-shi, Yamagata | N38-53-46.5 | E139-56-26.3 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-10 ② | 233624 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 山形県酒田市橋橋 | Narahashi, Sakata-shi, Yamagata | N38-53-46.5 | E139-56-26.3 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-10 ③ | 233625 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 山形県酒田市橋橋 | Narahashi, Sakata-shi, Yamagata | N38-53-46.5 | E139-56-26.3 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-11 | 233177 | <i>Glycine soja</i> | wild | 山形県東田川郡庄内町余目堤興屋 | Amarume, Shounai-machi, Higashi Tagawa-gun, Yamagata | N38-51-49.6 | E139-56-39.1 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-12 | 233178 | <i>Glycine soja</i> | wild | 山形県酒田市山寺 | Yamadera, Sakata-shi, Yamagata | N38-49-49.7 | E139-58-16.7 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-13 | 233179 | <i>Glycine soja</i> | wild | 山形県鶴岡市黒川 | Kurokawa, Tsuruoka-shi, Yamagata | N38-40-30.4 | E139-51-45.9 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-14 | 233180 | <i>Glycine soja</i> | wild | 山形県鶴岡市羽黒町松尾 | Matsuo, Haguro-machi, Tsuruoka-shi, Yamagata | N38-42-47.3 | E139-51-38.7 |
| 2007/9/24 | COL/Yamagata/2007/NIAS/CED2007-15 | 233181 | <i>Glycine soja</i> | wild | 山形県鶴岡市文下 | Houdashi, Tsuruoka-shi, Yamagata | N38-46-16.4 | E139-50-47.6 |
| 2007/10/22 | COL/Hyogo/2007/NIAS/CED2007Hyogo16-A | 233134 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 兵庫県美方郡新温泉町戸田 | Heda, Shin-Onsen-cho, Mikata-gun, Hyogo | N35-37-2.34 | E134-28-11.1 |
| 2007/10/22 | COL/Hyogo/2007/NIAS/CED2007Hyogo16-B | 233135 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 兵庫県美方郡新温泉町戸田 | Heda, Shin-Onsen-cho, Mikata-gun, Hyogo | N35-37-2.34 | E134-28-11.1 |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto17-A | 233136 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 京都府京丹後市久美浜町甲山地区 | Kouyama, Kumihama, Kyotango, Kyoto | N35-36-29.7 | E134-55-24.72 |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto17-B | 233137 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 京都府京丹後市久美浜町甲山地区 | Kouyama, Kumihama, Kyotango, Kyoto | N35-36-29.7 | E134-55-24.72 |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto17-C | 233138 | <i>Glycine max</i> | cultivated | 京都府京丹後市久美浜町甲山地区 | Kouyama, Kumihama, Kyotango, Kyoto | N35-36-29.7 | E134-55-24.72 |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto17-D | 233139 | <i>Glycine soja</i> | wild | 京都府京丹後市久美浜町甲山地区 | Kabutoyama, Kumihama, Kyoto | N35-36-29.7 | E134-55-24.72 |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto18-A | 233140 | <i>Glycine max</i> | cultivated | 京都府京丹後市久美浜町閑地区 | Seki, Kumihama, Kyotango, Kyoto | | |
| 2007/10/22 | COL/Kyoto/2007/NIAS/CED2007Kyoto18-B | 233141 | <i>Glycine soja</i> | wild | 京都府京丹後市久美浜町閑地区 | Seki, Kumihama, Kyotango, Kyoto | | |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo19-A | 233142 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 兵庫県豊岡市日高町藤井, 大目橋横 | R1, Ohme bridge, Yashiro river, Fujii, Hidaka, Toyooka, Hyogo | N35-29-37.5 | E134-46-58.5 |

| Altitude (m) | Habitat | Shading | Disturbance | Population size | Growth stage | Soil | Seed | Herbarium | Nodule | Remarks |
|--------------|-----------------------------------|---------|-------------|---|--|--------|------|-----------|--------|---|
| 5 | road side | open | high | 15 x 20m | flowering -mature | - | yes | no | no | unused land (road side) |
| 3 | river bank | open | medium | 50 x 100m | mature | sandy | yes | no | no | river bank of Akagawa, high density |
| 1 | river bank | open | medium | 40 x 200m | mature | - | yes | no | no | river bank of Mogamigawa, narrow leaf, an umbrella like pods setting type individual found here |
| 3 | road side | open | medium | 10 x 30m | mature | - | yes | no | no | unused land (road side) |
| 2 | beside railroad | open | high | 1 x 200m | mature | gravel | yes | no | no | unused land (road side) |
| 2 | hilly open place | open | medium | 15 x 40m | mature | - | yes | no | no | fallow field (margin) |
| 10 | river bank | open | medium | 10 x 10m | mature | - | yes | no | no | |
| 2 | river bank | open | medium | 3 x 100m | mature | - | yes | no | no | normal seed size, a plant with intermediate morphology between wild and cultivated found here |
| 2 | river bank | open | medium | 3 x 100m | mature | - | yes | no | no | large seed size |
| 10 | river side | open | medium | 5 x 1m | mature - past mature | - | yes | no | no | weedy type |
| 10 | river side | open | medium | 5 x 3m | mature - past mature | - | yes | no | no | wild type |
| 10 | river side | open | medium | - | mature - past mature | - | yes | no | no | probably escaped from cultivation |
| 12 | river bank | open | medium | 3 x 3m | flowering -mature - past mature | - | yes | no | no | beside rice field, wild type |
| 12 | river bank | open | medium | 3 x 30m | flowering -mature - past mature | - | yes | no | no | weedy type |
| 12 | river bank | open | medium | 3 x 30m | flowering -mature - past mature | - | yes | no | no | weedy type |
| 8 | river bank | open | medium | 50 x 50m | mature | - | yes | no | no | beside Mogamigawa |
| 8 | river bank | open | medium | 2 x 100m | mature | - | yes | no | no | low density |
| 48 | river bank | open | medium | 3 x 3m | mature | - | yes | no | no | beside Akagawa |
| 48 | river bank | open | medium | 5 x 2m | mature | - | yes | no | no | beside Akagawa |
| 46 | river bank | open | medium | 10 x 3m | mature | - | yes | no | no | abandoned place beside Akagawa, umbrella like pods setting type individuals found here |
| 5 | farmer's field | open | medium | 10 x 5m | mature | - | yes | yes | no | |
| 5 | river bank | open | medium | 2 x 5m | mature - past maturity | - | yes | yes | yes | |
| 1 | farmer's field | open | high | - | mature | - | yes | no | no | c.v. Kyoto Dainagon |
| 1 | beside field | open | high | 10 x 1m | mature | - | yes | no | no | short pods |
| 1 | farmer's field | open | high | - | mature | - | yes | no | no | c.v. Shin Tanbaguro, flowering time: 8/5 ~ 1month, Soy 2 years - Rice 2 years rotation |
| 1 | beside field | open | high | 50 x 2m | mature | - | yes | no | no | |
| - | farmer's field | open | high | - | mature | - | yes | no | no | |
| - | beside field | open | high | - | mature | - | yes | no | no | |
| 18 | river bank bushes beside paddy | light | low | triangle shaped place, 750m ² (50m x 30m ÷ 2) | past maturity | - | yes | yes | yes | 10 ~ 12 seeds/pod, short pod |

Table 2 (Continued).

| Coll.Date | Coll. No. | JP No. | Species name | Status | Collection Site (in Japanese) | Vill., City. Pref. | Latitude | Longitude |
|------------|--|--------|--|------------|----------------------------------|---|--------------|---------------|
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo19-B | 233143 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 兵庫県豊岡市日高町藤井, 大目橋横 | R1, Ohme bridge, Yashiro river, Fujii, Hidaka, Toyooka, Hyogo | N35-29-37.5 | E134-46-58.5 |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo20-A | 233144 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 兵庫県豊岡市竹野町桑野本（県道135号） | R135, Kuwanomoto, Takeno, Toyooka, Hyogo | N35-33-28.74 | E134-40-51.42 |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo20-B | 233145 | <i>Glycine soja</i> | wild | 兵庫県豊岡市竹野町桑野本（県道135号） | R135, Kuwanomoto, Takeno, Toyooka, Hyogo | N35-33-28.74 | E134-40-51.42 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto21-A | 233146 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 京都府京丹後市峰山町新治, 三軒屋橋横 | Ninbari, Mineyama, Kyotango, Kyoto | N35-36-22.44 | E135-03-15.3 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto21-B | 233147 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 京都府京丹後市峰山町新治, 三軒屋橋横 | Ninbari, Mineyama, Kyotango, Kyoto | N35-36-22.44 | E135-03-15.3 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto21-C | 233148 | <i>Glycine soja</i> | wild | 京都府京丹後市峰山町新治, 三軒屋橋横 | Ninbari, Mineyama, Kyotango, Kyoto | N35-36-22.44 | E135-03-15.3 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto22-A | 233149 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 京都府京丹後市丹後町宮竹野沖田 | Miya, Tango, Kyotango, Kyoto | N35-43-5.28 | E135-06-12.96 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto22-B | 233150 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 京都府京丹後市丹後町宮竹野沖田 | Miya, Tango, Kyotango, Kyoto | N35-43-5.28 | E135-06-12.96 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto22-C | 233151 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 京都府京丹後市丹後町宮竹野沖田 | Miya, Tango, Kyotango, Kyoto | N35-43-5.28 | E135-06-12.96 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto22-D | 233152 | <i>Glycine soja</i> | wild | 京都府京丹後市丹後町宮竹野沖田 | Miya, Tango, Kyotango, Kyoto | N35-43-5.28 | E135-06-12.96 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto23-A | 233153 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 京都府福知山市大江町千原村, 県道55号沿い | Senbara, Oe, Fukuchiyama, Kyoto | N35-22-47.22 | E135-9-28.8 |
| 2007/10/23 | COL/Kyoto/2007/NIAS/CED2007Kyoto23-B | 233154 | <i>Glycine soja</i> | wild | 京都府福知山市大江町千原村, 県道55号沿い | Senbara, Oe, Fukuchiyama, Kyoto | N35-22-47.22 | E135-9-28.8 |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo24-A(black mottled seed) | 233155 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 兵庫県丹波市市島町中竹田, 安下橋横 | Nakatakeda, Ichijima, Tanba, Hyogo | N35-14-23.22 | E135-07-55.02 |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo24-A(pale brown seed) | 233621 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 兵庫県丹波市市島町中竹田, 安下橋横 | Nakatakeda, Ichijima, Tanba, Hyogo | N35-14-23.22 | E135-07-55.02 |
| 2007/10/23 | COL/Hyogo/2007/NIAS/CED2007Hyogo24-B | 233156 | <i>Vigna angularis</i> var. <i>angularis</i> | cultivated | 兵庫県丹波市市島町中竹田, 安下橋横 | Nakatakeda, Ichijima, Tanba, Hyogo | N35-14-23.22 | E135-07-55.02 |
| 2007/10/24 | COL/Oita/2007/NIAS/CED2007Oita25 | 233157 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 大分県日田市内河野, 柳又発電所横 | Uchikawano, Hita, Oita | N33-18-18.18 | E130-53-7.86 |
| 2007/10/24 | COL/Fukuoka/2007/NIAS/CED2007Fukuoka26 | 233158 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 福岡県朝倉市山見, 県道80号横 | Yamami, Asakura, Fukuoka | N33-27-0.72 | E130-42-5.64 |
| 2007/10/24 | COL/Fukuoka/2007/NIAS/CED2007Fukuoka27 | 233159 | <i>Glycine soja</i> | wild | 福岡県筑紫野市山家, 下西山バス停横採石場（県道200号） | Yamae, Chikushino, Fukuoka | N33-30-7.92 | E130-35-41.28 |
| 2007/10/24 | COL/Saga/2007/NIAS/CED2007Saga28-A | 233160 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 佐賀県三養基郡基山町園部, 県道137号二瀬川橋横採石場 | Sonobe, Kiyama, Miyaki-gun, Saga | N33-25-31.68 | E130-29-32.82 |
| 2007/10/24 | COL/Saga/2007/NIAS/CED2007Saga28-B | 233161 | <i>Glycine soja</i> | wild | 佐賀県三養基郡基山町園部, 県道138号二瀬川橋横採石場 | Sonobe, Kiyama, Miyaki-gun, Saga | N33-25-31.68 | E130-29-32.82 |
| 2007/10/24 | COL/Saga/2007/NIAS/CED2007Saga28-C | 233162 | <i>Vigna unguiculata</i> | cultivated | 佐賀県三養基郡基山町園部, 県道139号二瀬川橋横採石場 | Sonobe, Kiyama, Miyaki-gun, Saga | N33-25-31.68 | E130-29-32.82 |
| 2007/10/25 | COL/Saga/2007/NIAS/CED2007Saga29-A | 233163 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 佐賀県佐賀市嘉瀬町十五（六字）, 干拓地, 下水浄化センター近く | Kase, Saga, Saga | N33-12-6.84 | E130-15-22.14 |
| 2007/10/25 | COL/Saga/2007/NIAS/CED2007Saga29-B | 233164 | <i>Glycine soja</i> | wild | 佐賀県佐賀市嘉瀬町十五（六字）, 干拓地, 下水浄化センター近く | Kase, Saga, Saga | N33-12-19.02 | E130-15-31.2 |
| 2007/10/25 | COL/Saga/2007/NIAS/CED2007Saga29-C | 233165 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 佐賀県佐賀市嘉瀬町十五（六字）, 干拓地, 下水浄化センター近く | Kase, Saga, Saga | N33-12-19.02 | E130-15-31.2 |
| 2007/11/5 | COL/Nagano/2007/NIAS/CED2007N1 | 233182 | <i>Glycine soja</i> | wild | 長野県伊那市南箕輪村小黒川バーチングエリア横の金網 | Ogurogawa, Minami Minowa Vill., Ina City, Nagano | N35-50-27.0 | E137-55-54.3 |
| 2007/11/5 | COL/Nagano/2007/NIAS/CED2007N2 | 233183 | <i>Glycine soja</i> | wild | 長野県伊那市南箕輪村, 信州大学の圃場横 | Shinshu Univ., Minami Minowa Vill., Ina, Nagano | N35-51-40.9 | E137-56-20.2 |
| 2007/11/6 | COL/Nagano/2007/NIAS/CED2007N3 | 233184 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 長野県泰阜村三耕地, 福寿院下 | Sankouchi, Yasuoka Vill., Nagano | N35-23-29.6 | E137-51-03.9 |
| 2007/11/6 | COL/Nagano/2007/NIAS/CED2007N4 | 233185 | <i>Vigna angularis</i> var. <i>nipponensis</i> | wild | 長野県阿南町大下条, 赤い屋根の保育園の前 | Ohshima, Anan town, Nagano | N35-19-24.9 | E137-48-49.2 |
| 2007/11/6 | COL/Nagano/2007/NIAS/CED2007N5 | 233186 | <i>Glycine soja</i> | wild | 長野県下條村睦沢, 県道64号沿い土砂採集場付近 | Mutsuzawa, Shimojo-Vill., Nagano | N35-24-27.6 | E137-46-55.0 |
| 2007/11/6 | COL/Nagano/2007/NIAS/CED2007N6 | 233187 | <i>Glycine soja</i> | wild | 長野県飯田市中平 | Nakahira, Iida City, Nagano | N35-27-39.3 | E1137-45-06.4 |
| 2007/11/6 | COL/Nagano/2007/NIAS/CED2007N7 | 233188 | <i>Vigna angularis</i> var. <i>nipponensis</i> | weedy | 長野県飯田市中平 | Nakahira, Iida City, Nagano | N35-27-39.3 | E1137-45-06.4 |

| Altitude (m) | Habitat | Shading | Disturbance | Population size | Growth stage | Soil | Seed | Herbarium | Nodule | Remarks |
|--------------|--------------------------------|--------------|-------------|-----------------|------------------------------------|-------|------|-----------|--------|--|
| 18 | river bank bushes beside paddy | light | low | 5 x 2m | past maturity | - | yes | yes | yes | larger leaves, possibility of introgression |
| 138 | fallow field | open | med - low | sporadically | past maturity | - | yes | yes | yes | fallow field (margin) and beside road |
| 138 | fallow field | open | med - low | sporadically | mature | - | yes | yes | yes | fallow field (margin) and beside road |
| 36 | river side bushes | open | medium | 5 x 5m | past maturity | - | yes | no | no | ca. 10~11 seeds/pod |
| 36 | river side bushes | open | medium | | - | - | yes | no | no | |
| 36 | river side bushes | open | medium | 2 x 2m | mature | - | yes | no | no | |
| 3 | beside road | open | high | 5 x 3m | past maturity | - | yes | no | yes | |
| 3 | cultivated field | open | high | | - | - | yes | no | no | white pod |
| 3 | cultivated field | open | high | | - | - | yes | no | no | brown pod, could be introgressed individual |
| 3 | river bank | open | high | 200 x 1m | mature | - | yes | no | yes | |
| 6 | river side | open | high | 10 x 1m | mature | - | yes | no | no | beside Yura river (R55), pod 5 ~ 6 cm, 12 seeds |
| 6 | river side | open | high | 10 x 2m | mature | - | yes | no | no | beside Yura river (R55), |
| 46 | bushes | open | medium | 30 x 50m | mature | - | yes | yes | no | abandoned place, seeds are small, Green stem + purple stem mixed, Black mottled seeded samples separated from mixed sample (CED2007Hyogo24-A). |
| 46 | bushes | open | medium | 30 x 50m | mature | - | yes | yes | no | abandoned place, seeds are small, Green stem + purple stem mixed, Pale brown seeded samples separated from mixed sample (CED2007Hyogo24-A). |
| 46 | cultivated field | open | medium | 30 x 50m | mature | - | yes | yes | no | |
| 93 | road side slope | open | high | 5 x 2m | flowering - mature - past maturity | - | yes | yes | yes | slope 20 ~ 30°, edge of road to stone wall, bracts, bracteole heavily hairy, bracteole much longer than carlyx. Green stem. 10 seeds/pod. Plants germinated after weeding. Some plants showed semi-indeterminate growth. Area least once cleared, stem nearly erect. |
| 93 | terraced paddy field | light - open | none | - | mature | - | yes | yes | no | slope 80°, stone wall of terraced paddy field, stem nearly glabrous |
| 146 | grassland | open | high | - | mature - past maturity | sandy | yes | no | no | growing abundantly on the ground of sand digging place of mountain, narrow leaflets |
| 130 | grassland | open | high | a few plants | past maturity | sandy | yes | no | no | mountain gravel digging place |
| 130 | grassland | open | high | - | mature | sandy | yes | no | no | mountain gravel digging place |
| 130 | farmer's field | open | high | 5 x 10m | maturing | sandy | yes | no | no | farmer's field beside mountain gravel digging place |
| 0 | beside soybean field | open | high | several plants | flowering - mature | - | yes | yes | yes | slope: 10°, field (soybean) margin, bracteole long and hairy, secondary bract large and hairy |
| 0 | beside irrigation canal | open | high | 5 x 1m | flowering - mature | - | yes | no | no | slope: 10° |
| 0 | beside irrigation canal | open | high | several plants | flowering - mature | - | yes | no | no | beside drainage, larger leaves and longer pods compared with 29-A |
| 736 | grassland | open | medium | 5 x 1m | past maturity | - | yes | no | no | climbing on the fence of Ogurogawa Parking Area of Chuo Highway |
| 756 | beside field | open | medium | several plants | past maturity | - | yes | no | no | climbing on the fence of experimental fields in Shinshu University |
| 699 | grassland | light | medium | many plants | past maturity | clay | yes | yes | yes | slope: 0 ~ 30°, beside paddy (slope), large seeds, stem green |
| 528 | beside path in paddy field | light | high | many plants | past maturity | clay | yes | yes | no | paddy field, 10 ~ 12 seeds/pod, stem purple, lobed leaflet, rather short pod, stem nearly hairless |
| 518 | road side | light | high | several plants | past maturity | - | yes | no | no | |
| 598 | waste land beside road | light | medium | many plants | past maturity | - | yes | no | no | |
| 598 | waste land beside road | light | medium | several plants | past maturity | - | yes | yes | yes | stem green, erect growth habit |



Photo 1. A wild soybean individual (*Glycine soja*) with umbrella like arrangement of pods found at 2007-03 site (Sakata city, Yamagata prefecture).




Photo 3. An individual with larger leaves and pods found among wild soybean (*G. soja*) population (2007-08) at Sakata city, Yamagata prefecture.



Photo 5. Wild azuki bean (left) and wild soybean (right) sometimes grow sympatrically in Japan (2007-22 site, Miya, Kyotango city, Kyoto prefecture).



Photo 2. A wild soybean individual (*Glycine soja*) with umbrella like arrangement of pods and few lateral branches. Most of pods are on main stem.



Photo 4. Plant type of a naturally growing soybean individual with larger leaves and pods (2007-08 site). This type may be hybrid derivative.



Photo 6. Variation in seed color and size within a naturally growing weedy azuki bean population (2007-24 site, Nakatakeda, Tanba-Ichijima, Hyogo prefecture).



Photo 7. A road side population of wild soybean (*G. soja*) found at Shimojo village, Nagano prefecture (2007N5).



Photo 8. Erect plant type of weedy azuki bean found in waste land beside road (Nakahira, Iida city, Nagano prefecture, 2007N7).